

University of Groningen

The bacterial sec machinery

van der Sluis, Elize Ouwe

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Sluis, E. O. (2006). *The bacterial sec machinery: structure, dynamics & evolution*. [Thesis fully internal (DIV), University of Groningen]. [s.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

1. Akita, M., A. Shinaki, S. Matsuyama, and S. Mizushima. 1991. SecA, an essential component of the secretory machinery of *Escherichia coli*, exists as homodimer. *Biochem.Biophys.Res.Commun.* **174**:211-216.
2. Baars, L., A. J. Ytterberg, D. Drew, S. Wagner, C. Thilo, K. J. van Wijk, and J. W. de Gier. 2005. Defining the role of the *E. coli* chaperone SecB using comparative proteomics. *J.Biol.Chem.* **281**:10024-10034.
3. Baba, T., T. Taura, T. Shimoike, Y. Akiyama, T. Yoshihisa, and K. Ito. 1994. A cytoplasmic domain is important for the formation of a SecY-SecE translocator complex. *Proc.Natl.Acad.Sci.U.S.A.* **91**:4539-4543.
4. Baud, C., S. Karamanou, G. Sianidis, E. Vrontou, A. S. Politou, and A. Economou. 2002. Allosteric communication between signal peptides and the SecA protein DEAD motor ATPase domain. *J.Biol.Chem.* **277**:13724-13731.
5. Beckmann, R., D. Bubeck, R. Grassucci, P. Penczek, A. Verschoor, G. Blobel, and J. Frank. 1997. Alignment of conduits for the nascent polypeptide chain in the ribosome-Sec61 complex. *Science* **278**:2123-2126.
6. Beckmann, R., C. M. Spahn, N. Eswar, J. Helmers, P. A. Penczek, A. Sali, J. Frank, and G. Blobel. 2001. Architecture of the protein-conducting channel associated with the translating 80S ribosome. *Cell* **107**:361-372.
7. Benach, J., Y. T. Chou, J. J. Fak, A. Itkin, D. D. Nicolae, P. C. Smith, G. Wittrock, D. L. Floyd, C. M. Golsaz, L. M. Gierasch, and J. F. Hunt. 2003. Phospholipid-induced monomerization and signal-peptide-induced oligomerization of SecA. *J.Biol.Chem.* **278**:3628-3638.
8. Bensing, B. A., B. W. Gibson, and P. M. Sullam. 2004. The *Streptococcus gordonii* platelet binding protein GspB undergoes glycosylation independently of export. *J.Bacteriol.* **186**:638-645.
9. Bensing, B. A. and P. M. Sullam. 2002. An accessory *sec* locus of *Streptococcus gordonii* is required for export of the surface protein GspB and for normal levels of binding to human platelets. *Mol.Microbiol.* **44**:1081-1094.
10. Bensing, B. A., D. Takamatsu, and P. M. Sullam. 2005. Determinants of the streptococcal surface glycoprotein GspB that facilitate export by the accessory Sec system. *Mol.Microbiol.* **58**:1468-1481.
11. Bessonneau, P., V. Besson, I. Collinson, and F. Duong. 2002. The SecYEG preprotein translocation channel is a conformationally dynamic and dimeric structure. *EMBO J.* **21**:995-1003.
12. Beswick, V., F. Baleux, T. Huynh-Dinh, F. Kepes, J. M. Neumann, and A. Sanson. 1996. NMR conformational study of the cytoplasmic domain of the canine Sec61 γ protein from the protein translocation pore of the endoplasmic reticulum membrane. *Biochemistry* **35**:14717-14724.
13. Bost, S. and D. Belin. 1995. A new genetic selection identifies essential residues in SecG, a component of the *Escherichia coli* protein export machinery. *EMBO J.* **14**:4412-4421.
14. Bostina, M., B. Mohsin, W. Kuhlbrandt, and I. Collinson. 2005. Atomic model of the *E. coli* membrane-bound protein translocation complex SecYEG. *J.Mol.Biol.* **352**:1035-1043.
15. Braunstein, M., A. M. Brown, S. Kurtz, and W. R. Jacobs, Jr. 2001. Two nonredundant SecA homologues function in Mycobacteria. *J.Bacteriol.* **183**:6979-6990.
16. Braunstein, M., B. J. Espinosa, J. Chan, J. T. Belisle, and R. Jacobs. 2003. SecA2 functions in the secretion of superoxide dismutase A and in the virulence of *Mycobacterium tuberculosis*. *Mol.Microbiol.* **48**:453-464.

34. **Dapic, V. and D. Oliver.** 2000. Distinct membrane binding properties of N- and C-terminal domains of *Escherichia coli* SecA ATPase. *J.Biol.Chem.* **275**:25000-25007.
35. **De Keyzer, J., C. van der Does, and A. J. M. Driessen.** 2002. Kinetic analysis of the translocation of fluorescent precursor proteins into *Escherichia coli* membrane vesicles. *J.Biol.Chem.* **277**:46059-46065.
36. **De Keyzer, J., C. van der Does, and A. J. M. Driessen.** 2003. The bacterial translocase: a dynamic protein channel complex. *Cell Mol.Life Sci.* **60**:2034-2052.
37. **De Keyzer, J., C. van der Does, T. G. Kloosterman, and A. J. M. Driessen.** 2003. Direct demonstration of ATP-dependent release of SecA from a translocating preprotein by Surface Plasmon Resonance. *J.Biol.Chem.* **278**:29581-29586.
38. **De Keyzer, J., C. van der Does, J. Swaving, and A. J. M. Driessen.** 2002. The F286Y mutation of PrlA4 tempers the signal sequence suppressor phenotype by reducing the SecA binding affinity. *FEBS Lett.* **510**:17-21.
39. **De Keyzer, J., E. O. van der Sluis, R. E. Spelbrink, N. Nijstad, B. de Kruijff, N. Nouwen, C. van der Does, and A. J. M. Driessen.** 2005. Covalently dimerized SecA is functional in protein translocation. *J.Biol.Chem.* **280**:35255-35260.
40. **Dekker, C., B. de Kruijff, and P. Gros.** 2003. Crystal structure of SecB from *Escherichia coli*. *J.Struct.Biol.* **144**:313-319.
41. **Delepelaire, P. and C. Wandersman.** 1998. The SecB chaperone is involved in the secretion of the *Serratia marcescens* HasA protein through an ABC transporter. *EMBO J.* **17**:936-944.
42. **Delsuc, F., H. Brinkmann, and H. Philippe.** 2005. Phylogenomics and the reconstruction of the tree of life. *Nat.Rev.Genet.* **6**:361-375.
43. **Dempsey, B. R., A. Economou, S. D. Dunn, and B. H. Shilton.** 2002. The ATPase domain of SecA can form a tetramer in solution. *J.Mol.Biol.* **315**:831-843.
44. **Dempsey, B. R., M. Wrona, J. M. Moulin, G. B. Gloor, F. Jalilehvand, G. Lajoie, G. S. Shaw, and B. H. Shilton.** 2004. Solution NMR structure and X-ray absorption analysis of the C-terminal zinc-binding domain of the SecA ATPase. *Biochemistry.* **43** :9361-9371.
45. **Den Blaauwen, T., P. Fekkes, J. G. de Wit, W. Kuiper, and A. J. M. Driessen.** 1996. Domain interactions of the peripheral preprotein translocase subunit SecA. *Biochemistry* **35**:11994-12004.
46. **Ding, H., J. F. Hunt, I. Mukerji, and D. Oliver.** 2003. *Bacillus subtilis* SecA ATPase exists as an antiparallel dimer in solution. *Biochemistry* **42**:8729-8738.
47. **Ding, H., I. Mukerji, and D. Oliver.** 2001. Lipid and signal peptide-induced conformational changes within the C-domain of *Escherichia coli* SecA protein. *Biochemistry* **40**:1835-1843.
48. **Ding, H., I. Mukerji, and D. Oliver.** 2003. Nucleotide and phospholipid-dependent control of PPXD and C-domain association for SecA ATPase. *Biochemistry* **42**:13468-13475.
49. **Driessen, A. J. M.** 1993. SecA, the peripheral subunit of the *Escherichia coli* precursor protein translocase, is functional as a dimer. *Biochemistry* **32**:13190-13197.
50. **Driessen, A. J. M., E. H. Manting, and C. van der Does.** 2001. The structural basis of protein targeting and translocation in bacteria. *Nat.Struct.Biol.* **8**:492-498.
51. **Duong, F.** 2003. Binding, activation and dissociation of the dimeric SecA ATPase at the dimeric SecYEG translocase. *EMBO J.* **22**:4375-4384.

52. **Duong, F. and W. Wickner.** 1997. Distinct catalytic roles of the SecYE, SecG and SecDFyajC subunits of preprotein translocase holoenzyme. *EMBO J.* **16**:2756-2768.
53. **Duong, F. and W. Wickner.** 1997. The SecDFyajC domain of preprotein translocase controls preprotein movement by regulating SecA membrane cycling. *EMBO J.* **16**:4871-4879.
54. **Economou, A., J. A. Pogliano, J. Beckwith, D. B. Oliver, and W. Wickner.** 1995. SecA membrane cycling at SecYEG is driven by distinct ATP binding and hydrolysis events and is regulated by SecD and SecF. *Cell* **83**:1171-1181.
55. **Economou, A. and W. Wickner.** 1994. SecA promotes preprotein translocation by undergoing ATP-driven cycles of membrane insertion and deinsertion. *Cell* **78**:835-843.
56. **Eichler, J., K. Rinard, and W. Wickner.** 1998. Endogenous SecA catalyzes preprotein translocation at SecYEG. *J.Biol.Chem.* **273**:21675-21681.
57. **Fekkes, P., J. G. de Wit, A. Boorsma, R. H. Friesen, and A. J. M. Driessen.** 1999. Zinc stabilizes the SecB binding site of SecA. *Biochemistry* **38**:5111-5116.
58. **Fekkes, P., J. G. de Wit, J. P. van der Wolk, H. H. Kimsey, C. A. Kumamoto, and A. J. M. Driessen.** 1998. Preprotein transfer to the *Escherichia coli* translocase requires the co-operative binding of SecB and the signal sequence to SecA. *Mol.Microbiol.* **29**:1179-1190.
59. **Fekkes, P., C. van der Does, and A. J. M. Driessen.** 1997. The molecular chaperone SecB is released from the carboxy-terminus of SecA during initiation of precursor protein translocation. *EMBO J.* **16**:6105-6113.
60. **Finke, K., K. Plath, S. Panzner, S. Prehn, T. A. Rapoport, E. Hartmann, and T. Sommer.** 1996. A second trimeric complex containing homologs of the Sec61p complex functions in protein transport across the ER membrane of *S. cerevisiae*. *EMBO J.* **15**:1482-1494.
61. **Flower, A. M.** 2001. SecG function and phospholipid metabolism in *Escherichia coli*. *J.Bacteriol.* **183**:2006-2012.
62. **Flower, A. M., L. L. Hines, and P. L. Pfennig.** 2000. SecG is an auxiliary component of the protein export apparatus of *Escherichia coli*. *Mol.Gen.Genet.* **263**:131-136.
63. **Flower, A. M., R. S. Osborne, and T. J. Silhavy.** 1995. The allele-specific synthetic lethality of *prlA-prlG* double mutants predicts interactive domains of SecY and SecE. *EMBO J.* **14**:884-893.
64. **Frank, R. and H. Overwin.** 1996. SPOT synthesis. Epitope analysis with arrays of synthetic peptides prepared on cellulose membranes. *Methods Mol.Biol.* **66**:149-69.:149-169.
65. **Friedrich, T. and H. Weiss.** 1997. Modular evolution of the respiratory NADH:ubiquinone oxidoreductase and the origin of its modules. *J.Theor.Biol.* **187**:529-540.
66. **Gabaldon, T., D. Rainey, and M. A. Huynen.** 2005. Tracing the evolution of a large protein complex in the eukaryotes, NADH:ubiquinone oxidoreductase (Complex I). *J.Mol.Biol.* **348**:857-870.
67. **Gannon, P. M., P. Li, and C. A. Kumamoto.** 1989. The mature portion of *Escherichia coli* maltose-binding protein (MBP) determines the dependence of MBP on SecB for export. *J.Bacteriol.* **171**:813-818.
68. **Groves, M. R., A. Mant, A. Kuhn, J. Koch, S. Dubel, C. Robinson, and I. Sinning.** 2001. Functional characterization of recombinant chloroplast signal recognition particle. *J.Biol.Chem.* **276**:27778-27786.

69. **Hanada, M., K. Nishiyama, S. Mizushima, and H. Tokuda.** 1994. Reconstitution of an efficient protein translocation machinery comprising SecA and the three membrane proteins, SecY, SecE, and SecG (p12). *J.Biol.Chem.* **269**:23625-23631.
70. **Hanein, D., K. E. Matlack, B. Jungnickel, K. Plath, K. U. Kalies, K. R. Miller, T. A. Rapoport, and C. W. Akey .** 1996. Oligomeric rings of the Sec61p complex induced by ligands required for protein translocation. *Cell* **87**:721-732.
71. **Harris, C. R. and T. J. Silhavy.** 1999. Mapping an interface of SecY (PrlA) and SecE (PrlG) by using synthetic phenotypes and *in vivo* cross-linking. *J.Bacteriol.* **181**:3438-3444.
72. **Hartl, F. U., S. Lecker, E. Schiebel, J. P. Hendrick, and W. Wickner.** 1990. The binding cascade of SecB to SecA to SecY/E mediates preprotein targeting to the *E. coli* plasma membrane. *Cell* **63**:269-279.
73. **Homma, T., T. Yoshihisa, and K. Ito.** 1997. Subunit interactions in the *Escherichia coli* protein translocase: SecE and SecG associate independently with SecY. *FEBS Lett.* **408**:11-15.
74. **Hunt, J. F., S. Weinkauf, L. Henry, J. J. Fak, P. McNicholas, D. B. Oliver, and J. Deisenhofer.** 2002. Nucleotide control of interdomain interactions in the conformational reaction cycle of SecA. *Science* **297**:2018-2026.
75. **Huynen, M. A., T. Gabaldon, and B. Snel.** 2005. Variation and evolution of biomolecular systems: searching for functional relevance. *FEBS Lett.* **579**:1839-1845.
76. **Jia, L., M. Dienhart, M. Schramp, M. McCauley, K. Hell, and R. A. Stuart.** 2003. Yeast Oxa1 interacts with mitochondrial ribosomes: the importance of the C-terminal region of Oxa1. *EMBO J.* **22**:6438-6447.
77. **Jilaveanu, L. B. and D. Oliver .** 2006. SecA dimer cross-linked at its subunit interface is functional for protein translocation. *J.Bacteriol.* **188**:335-338.
78. **Jilaveanu, L. B., C. R. Zito, and D. Oliver.** 2005. Dimeric SecA is essential for protein translocation. *Proc.Natl.Acad.Sci.U.S.A.* **102**:7511-7516.
79. **Joly, J. C. and W. T. Wickner.** 1993. The SecA and SecY subunits of translocase are the nearest neighbors of a translocating preprotein, shielding it from phospholipids. *EMBO J.* **12**:255-263.
80. **Kalies, K. U., D. Gorlich, and T. A. Rapoport.** 1994. Binding of ribosomes to the rough endoplasmic reticulum mediated by the Sec61p-complex. *J.Cell Biol.* **126**:925-934.
81. **Kalies, K. U., T. A. Rapoport, and E. Hartmann.** 1998. The β -subunit of the Sec61 complex facilitates cotranslational protein transport and interacts with the signal peptidase during translocation. *J.Cell Biol.* **141**:887-894.
82. **Karamanou, S., G. Sianidis, G. Gouridis, C. Pozidis, Y. Papanikolau, E. Papanikou, and A. Economou.** 2005. *Escherichia coli* SecA truncated at its termini is functional and dimeric. *FEBS Lett.* **579**:1267-1271.
83. **Karamyshev, A. L. and A. E. Johnson.** 2005. Selective SecA association with signal sequences in ribosome-bound nascent chains: a potential role for SecA in ribosome targeting to the bacterial membrane. *J.Biol.Chem.* **280**:37930-37940.
84. **Kato, Y., K. Nishiyama, and H. Tokuda.** 2003. Depletion of SecDF-YajC causes a decrease in the level of SecG: implication for their functional interaction. *FEBS Lett.* **550**:114-118.
85. **Kaufmann, A., E. H. Manting, A. K. J. Veenendaal, A. J. M. Driessen, and C. van der Does.** 1999. Cysteine-directed cross-linking demonstrates that helix 3 of SecE is close to helix 2 of SecY and helix 3 of a neighboring SecE. *Biochemistry* **38**:9115-9125.

86. **Kimsey, H. H., M. D. Dagarag, and C. A. Kumamoto.** 1995. Diverse effects of mutation on the activity of the *Escherichia coli* export chaperone SecB. *J.Biol.Chem.* **270**:22831-22835.
87. **Kimura, E., M. Akita, S. Matsuyama, and S. Mizushima.** 1991. Determination of a region in SecA that interacts with presecretory proteins in *Escherichia coli*. *J.Biol.Chem.* **266**:6600-6606.
88. **Kinch, L. N., M. H. Saier, Jr., and N. V. Grishin.** 2002. Sec61 β -a component of the archaeal protein secretory system. *Trends Biochem.Sci.* **27**:170-171.
89. **Knoblauch, N. T., S. Rudiger, H. J. Schonfeld, A. J. M. Driessen, J. Schneider-Mergener, and B. Bukau.** 1999. Substrate specificity of the SecB chaperone. *J.Biol.Chem.* **274**:34219-34225.
90. **Koch, J. and M. Mahler.** 2002. Peptide arrays on membrane supports. Springer Verlag, Heidelberg, Germany .
91. **Koonin, E. V. and A. E. Gorbalenya.** 1992. Autogenous translation regulation by *Escherichia coli* ATPase SecA may be mediated by an intrinsic RNA helicase activity of this protein. *FEBS Lett.* **298**:6-8.
92. **Kumamoto, C. A. and J. Beckwith.** 1985. Evidence for specificity at an early step in protein export in *Escherichia coli*. *J.Bacteriol.* **163**:267-274.
93. **Kunin, V., L. Goldovsky, N. Darzentas, and C. A. Ouzounis.** 2005. The net of life: reconstructing the microbial phylogenetic network. *Genome Res.* **15**:954-959.
94. **Larsson, P., P. C. Oyston, P. Chain, M. C. Chu, M. Duffield, H. H. Fuxelius, E. Garcia, G. Halltorp, D. Johansson, K. E. Isherwood, P. D. Karp, E. Larsson, Y. Liu, S. Michell, J. Prior, R. Prior, S. Malfatti, A. Sjostedt, K. Svensson, N. Thompson, L. Vergez, J. K. Wagg, B. W. Wren, L. E. Lindler, S. G. Andersson, M. Forsman, and R. W. Titball.** 2005. The complete genome sequence of *Francisella tularensis*, the causative agent of tularemia. *Nat.Genet.* **37**:153-159.
95. **Lecker, S. H., A. J. M. Driessen, and W. Wickner.** 1990. ProOmpA contains secondary and tertiary structure prior to translocation and is shielded from aggregation by association with SecB protein. *EMBO J.* **9**:2309-2314.
96. **Lenz, L. L., S. Mohammadi, A. Geissler, and D. A. Portnoy.** 2003. SecA2-dependent secretion of autolytic enzymes promotes *Listeria monocytogenes* pathogenesis. *Proc.Natl.Acad.Sci.U.S.A* **100**:12432-12437.
97. **Lenz, L. L. and D. A. Portnoy.** 2002. Identification of a second *Listeria* secA gene associated with protein secretion and the rough phenotype. *Mol.Microbiol.* **45**:1043-1056.
98. **Lerat, E., V. Daubin, and N. A. Moran.** 2003. From gene trees to organismal phylogeny in prokaryotes: the case of the γ -Proteobacteria. *PLoS.Biol.* **1**:E19.
99. **Liebke, H. H.** 1987. Multiple SecA protein isoforms in *Escherichia coli*. *J.Bacteriol.* **169**:1174-1181.
100. **Lill, R., K. Cunningham, L. A. Brundage, K. Ito, D. Oliver, and W. Wickner.** 1989. SecA protein hydrolyzes ATP and is an essential component of the protein translocation ATPase of *Escherichia coli*. *EMBO J.* **8**:961-966.
101. **Luirink, J. and I. Sinning.** 2004. SRP-mediated protein targeting: structure and function revisited. *Biochim.Biophys.Acta.* **1694**:17-35.
102. **Machata, S., T. Hain, M. Rohde, and T. Chakraborty.** 2005. Simultaneous deficiency of both MurA and p60 proteins generates a rough phenotype in *Listeria monocytogenes*. *J.Bacteriol.* **187**:8385-8394.

-
103. **Manting, E. H., C. van der Does, and A. J. M. Driessen.** 1997. *In vivo* cross-linking of the SecA and SecY subunits of the *Escherichia coli* preprotein translocase. *J.Bacteriol.* **179**:5699-5704.
 104. **Manting, E. H., C. van der Does, H. Remigy, A. Engel, and A. J. M. Driessen.** 2000. SecYEG assembles into a tetramer to form the active protein translocation channel. *EMBO J.* **19**:852-861.
 105. **Mason, N., L. F. Ciuffo, and J. D. Brown.** 2000. Elongation arrest is a physiologically important function of signal recognition particle. *EMBO J.* **19**:4164-4174.
 106. **Mathiesen, C. and C. Hagerhall.** 2003. The 'antiporter module' of respiratory chain complex I includes the MrpC/NuoK subunit -- a revision of the modular evolution scheme. *FEBS Lett.* **549**:7-13.
 107. **Matousek, W. M. and A. T. Alexandrescu.** 2004. NMR structure of the C-terminal domain of SecA in the free state. *Biochim.Biophys.Acta.* **1702**:163-171.
 108. **Matsumoto, G., H. Mori, and K. Ito.** 1998. Roles of SecG in ATP- and SecA-dependent protein translocation. *Proc.Natl.Acad.Sci.U.S.A.* **95**:13567-13572.
 109. **Matsumoto, G., H. Nakatogawa, H. Mori, and K. Ito.** 2000. Genetic dissection of SecA: suppressor mutations against the SecY205 translocase defect. *Genes Cells* **5**:991-999.
 110. **Matsuyama, S., J. Akimaru, and S. Mizushima.** 1990. SecE-dependent overproduction of SecY in *Escherichia coli*. Evidence for interaction between two components of the secretory machinery. *FEBS Lett.* **269**:96-100.
 111. **Matsuyama, S., E. Kimura, and S. Mizushima.** 1990. Complementation of two overlapping fragments of SecA, a protein translocation ATPase of *Escherichia coli*, allows ATP binding to its amino-terminal region. *J.Biol.Chem.* **265**:8760-8765.
 112. **Menetret, J. F., R. S. Hegde, S. U. Heinrich, P. Chandramouli, S. J. Ludtke, T. A. Rapoport, and C. W. Akey.** 2005. Architecture of the ribosome-channel complex derived from native membranes. *J.Mol.Biol.* **348**:445-457.
 113. **Menetret, J. F., A. Neuhofer, D. G. Morgan, K. Plath, M. Radermacher, T. A. Rapoport, and C. W. Akey.** 2000. The structure of ribosome-channel complexes engaged in protein translocation. *Mol.Cell* **6**:1219-1232.
 114. **Meyer, T. H., J. F. Ménétret, R. Breitling, K. R. Miller, C. W. Akey, and T. A. Rapoport.** 1999. The Bacterial SecYEG translocation complex forms channel-like structures similar to those of the eukaryotic Sec61p Complex. *J.Mol.Biol.* **285**:1789-1800.
 115. **Mitra, K., C. Schaffitzel, T. Shaikh, F. Tama, S. Jenni, C. L. Brooks, III, N. Ban, and J. Frank.** 2005. Structure of the *E. coli* protein-conducting channel bound to a translating ribosome. *Nature.* **438**:318-324.
 116. **Morgan, D. G., J. F. Menetret, A. Neuhofer, T. A. Rapoport, and C. W. Akey.** 2002. Structure of the mammalian ribosome-channel complex at 17Å resolution. *J.Mol.Biol.* **324**:871-886.
 117. **Morgan, D. G., J. F. Menetret, M. Radermacher, A. Neuhofer, I. V. Akey, T. A. Rapoport, and C. W. Akey.** 2000. A comparison of the yeast and rabbit 80 S ribosome reveals the topology of the nascent chain exit tunnel, inter-subunit bridges and mammalian rRNA expansion segments. *J.Mol.Biol.* **301**:301-321.
 118. **Mori, H. and K. Ito.** 2001. An essential amino acid residue in the protein translocation channel revealed by targeted random mutagenesis of SecY. *Proc.Natl.Acad.Sci.U.S.A* **98**:5128-5133.
 119. **Mori, H. and K. Ito.** 2001. The Sec protein-translocation pathway. *Trends Microbiol.* **9**:494-500.

120. **Mori, H. and K. Ito.** 2003. Biochemical characterization of a mutationally altered protein translocase: proton motive force stimulation of the initiation phase of translocation. *J.Bacteriol.* **185**:405-412.
121. **Mori, H., N. Shimokawa, Y. Satoh, and K. Ito.** 2004. Mutational analysis of transmembrane regions 3 and 4 of SecY, a central component of protein translocase. *J.Bacteriol.* **186**:3960-3969.
122. **Mori, H., H. Sugiyama, M. Yamanaka, K. Sato, M. Tagaya, and S. Mizushima.** 1998. Amino-terminal region of SecA is involved in the function of SecE for protein translocation into *Escherichia coli* membrane vesicles. *J.Biochem.(Tokyo)* **124**:122-129.
123. **Mori, H., T. Tsukazaki, R. Masui, S. Kuramitsu, S. Yokoyama, A. E. Johnson, Y. Kimura, Y. Akiyama, and K. Ito .** 2003. Fluorescence resonance energy transfer analysis of protein translocase. SecYE from *Thermus thermophilus* HB8 forms a constitutive oligomer in membranes. *J.Biol.Chem.* **278**:14257-14264.
124. **Musial-Siwiek, M., S. L. Rusch, and D. A. Kendall.** 2005. Probing the affinity of SecA for signal peptide in different environments. *Biochemistry.* **44**:13987-13996.
125. **Nagamori, S., K. Nishiyama, and H. Tokuda.** 2000. Two SecE molecules present in a single protein translocation machinery are functional even after crosslinking. *J.Biochem.* **128**:129-137.
126. **Nagamori, S., K. Nishiyama, and H. Tokuda.** 2002. Membrane topology inversion of SecE detected by labeling with a membrane-impermeable sulfhydryl reagent that causes a close association of SecE with SecA. *J.Biochem.(Tokyo)* **132**:629-634.
127. **Nakatogawa, H. and K. Ito.** 2001. Secretion monitor, SecM, undergoes self-translation arrest in the cytosol. *Mol.Cell* **7**:185-192.
128. **Nakatogawa, H. and K. Ito.** 2002. The ribosomal exit tunnel functions as a discriminating gate. *Cell* **108**:629-636.
129. **Nakatogawa, H., A. Murakami, H. Mori, and K. Ito.** 2005. SecM facilitates translocase function of SecA by localizing its biosynthesis. *Genes Dev.* **19**:436-444.
130. **Neumann-Haefelin, C., U. Schafer, M. Muller, and H. G. Koch.** 2000. SRP-dependent co-translational targeting and SecA-dependent translocation analyzed as individual steps in the export of a bacterial protein. *EMBO J.* **19**:6419-6426.
131. **Nishiyama, K., M. Hanada, and H. Tokuda.** 1994. Disruption of the gene encoding p12 (SecE) reveals the direct involvement and important function of SecE in the protein translocation of *Escherichia coli* at low temperature. *EMBO J.* **13**:3272-3277.
132. **Nishiyama, K., S. Mizushima, and H. Tokuda.** 1993. A novel membrane protein involved in protein translocation across the cytoplasmic membrane of *Escherichia coli*. *EMBO J.* **12**:3409-3415.
133. **Nishiyama, K., T. Suzuki, and H. Tokuda.** 1996. Inversion of the membrane topology of SecE coupled with SecA-dependent preprotein translocation. *Cell* **85**:71-81.
134. **Nouwen, N. and A. J. M. Driessen.** 2002. SecDFyajC forms a heterotetrameric complex with YidC. *Mol.Microbiol.* **44**:1397-1405.
135. **Nouwen, N., M. van der Laan, and A. J. M. Driessen.** 2001. SecDFyajC is not required for the maintenance of the proton motive force. *FEBS Lett.* **508**:103-106.
136. **Or, E., D. Boyd, S. Gon, J. Beckwith, and T. Rapoport.** 2005. The bacterial ATPase SecA functions as a monomer in protein translocation. *J.Biol.Chem.* **280**:9097-9105.
137. **Or, E., A. Navon, and T. Rapoport.** 2002. Dissociation of the dimeric SecA ATPase during protein translocation across the bacterial membrane. *EMBO J.* **21**:4470-4479.

138. **Osborne, A. R., W. M. Clemons, Jr., and T. A. Rapoport.** 2004. A large conformational change of the translocation ATPase SecA. *Proc.Natl.Acad.Sci.U.S.A.* **101**:10937-10942.
139. **Osborne, A. R., T. A. Rapoport, and B. van den Berg.** 2005. Protein translocation by the Sec61/SecY channel. *Annu.Rev.Cell Dev.Biol.* **21**:529-50.:529-550.
140. **Overath, P., M. Brenner, T. Gulik-Krzywicki, E. Shechter, and L. Letellier.** 1975. Lipid phase transitions in cytoplasmic and outer membranes of *Escherichia coli*. *Biochim.Biophys.Acta.* **389**:358-369.
141. **Papanikou, E., S. Karamanou, C. Baud, M. Frank, G. Sianidis, D. Keramisanou, C. G. Kalodimos, A. Kuhn, and A. Economou.** 2005. Identification of the preprotein binding domain of SecA. *J.Biol.Chem.* **280**:43209-43217.
142. **Plath, K., W. Mothes, B. M. Wilkinson, C. J. Stirling, and T. A. Rapoport.** 1998. Signal sequence recognition in posttranslational protein transport across the yeast ER membrane. *Cell* **94**:795-807.
143. **Plath, K., B. M. Wilkinson, C. J. Stirling, and T. A. Rapoport.** 2004. Interactions between Sec complex and prepro-alpha-factor during posttranslational protein transport into the endoplasmic reticulum. *Mol.Biol.Cell.* **15**:1-10.
144. **Pohlschroder, M., E. Hartmann, N. J. Hand, K. Dilks, and A. Haddad.** 2005. Diversity and evolution of protein translocation. *Annu.Rev.Microbiol.* **59**:91-111.:91-111.
145. **Pohlschroder, M., C. Murphy, and J. Beckwith.** 1996. *In vivo* analyses of interactions between SecE and SecY, core components of the *Escherichia coli* protein translocation machinery. *J.Biol.Chem.* **271**:19908-19914.
146. **Pohlschroder, M., W. A. Prinz, E. Hartmann, and J. Beckwith.** 1997. Protein translocation in the three domains of life: variations on a theme. *Cell* **91**:563-566.
147. **Prinz, A., C. Behrens, T. A. Rapoport, E. Hartmann, and K. U. Kalies.** 2000. Evolutionarily conserved binding of ribosomes to the translocation channel via the large ribosomal RNA. *EMBO J.* **19**:1900-1906.
148. **Raden, D., W. Song, and R. Gilmore.** 2000. Role of the cytoplasmic segments of Sec61 α in the ribosome-binding and translocation-promoting activities of the Sec61 complex. *J.Cell Biol.* **150**:53-64.
149. **Rain, J. C., L. Selig, H. De Reuse, V. Battaglia, C. Reverdy, S. Simon, G. Lenzen, F. Petel, J. Wojcik, V. Schachter, Y. Chemama, A. Labigne, and P. Legrain.** 2001. The protein-protein interaction map of *Helicobacter pylori*. *Nature.* **409**:211-215.
150. **Rajapandi, T., K. M. Dolan, and D. B. Oliver.** 1991. The first gene in the *Escherichia coli* *secA* operon, gene X, encodes a nonessential secretory protein. *J.Bacteriol.* **173**:7092-7097.
151. **Rajapandi, T. and D. Oliver.** 1994. Carboxy-terminal region of *Escherichia coli* SecA ATPase is important to promote its protein translocation activity *in vivo*. *Biochem.Biophys.Res.Comm.* **200**:1477-1483.
152. **Randall, L. L., J. M. Crane, A. A. Lilly, G. Liu, C. Mao, C. N. Patel, and S. J. Hardy.** 2005. Asymmetric binding between SecA and SecB two symmetric proteins: implications for function in export. *J.Mol.Biol.* **348**:479-489.
153. **Rapiejko, P. J. and R. Gilmore.** 1997. Empty site forms of the SRP54 and SR α GTPases mediate targeting of ribosome-nascent chain complexes to the endoplasmic reticulum. *Cell* **89**:703-713.
154. **Samuelson, J. C., F. Jiang, L. Yi, M. Chen, J. W. de Gier, A. Kuhn, and R. E. Dalbey.** 2001. Function of YidC for the insertion of M13 procoat protein in *Escherichia coli*: translocation of

- mutants that show differences in their membrane potential dependence and Sec requirement. *J.Biol.Chem.* **276** :34847-34852.
155. Sato, K., H. Mori, M. Yoshida, M. Tagaya, and S. Mizushima. 1997. *In vitro* analysis of the stop-transfer process during translocation across the cytoplasmic membrane of *Escherichia coli*. *J.Biol.Chem.* **272**:20082-20087.
 156. Satoh, Y., G. Matsumoto, H. Mori, and K. Ito. 2003. Nearest neighbor analysis of the SecYEG complex. 1. Identification of a SecY-SecG interface. *Biochemistry* **42**:7434-7441.
 157. Satoh, Y., H. Mori, and K. Ito. 2003. Nearest neighbor analysis of the SecYEG complex. 2. Identification of a SecY-SecE cytosolic interface. *Biochemistry* **42**:7442-7447.
 158. Schatz, P. J., K. L. Bieker, K. M. Ottemann, T. J. Silhavy, and J. Beckwith. 1991. One of three transmembrane stretches is sufficient for the functioning of the SecE protein, a membrane component of the *E. coli* secretion machinery. *EMBO J.* **10**:1749-1757.
 159. Schatz, P. J., P. D. Riggs, A. Jacq, M. J. Fath, and J. Beckwith. 1989. The *secE* gene encodes an integral membrane protein required for protein export in *Escherichia coli*. *Genes Dev.* **3**:1035-1044.
 160. Scheuring, J., N. Braun, L. Nothdurft, M. Stumpf, A. K. Veenendaal, S. Kol, C. van der Does, A. J. M. Driessen, and S. Weinkauf. 2005. The oligomeric distribution of SecYEG is altered by SecA and translocation ligands. *J.Mol.Biol.* **354**:258-271.
 161. Schiebel, E., A. J. M. Driessen, F. U. Hartl, and W. Wickner. 1991. $\Delta\mu\text{H}^+$ and ATP function at different steps of the catalytic cycle of preprotein translocase. *Cell* **64**:927-939.
 162. Schmidt, M., H. Ding, V. Ramamurthy, I. Mukerji, and D. Oliver. 2000. Nucleotide binding activity of SecA homodimer is conformationally regulated by temperature and altered by *prlD* and *azl* mutations. *J.Biol.Chem.* **275**:15440-15448.
 163. Schuwirth, B. S., M. A. Borovinskaya, C. W. Hau, W. Zhang, A. Vila-Sanjurjo, J. M. Holton, and J. H. Cate. 2005. Structures of the bacterial ribosome at 3.5 Å resolution. *Science*. **310**:827-834.
 164. Scotti, P. A., M. L. Urbanus, J. Brunner, J. W. de Gier, G. von Heijne, C. van der Does, A. J. M. Driessen, B. Oudega, and J. Luirink. 2000. YidC, the *Escherichia coli* homologue of mitochondrial Oxa1p, is a component of the Sec translocase. *EMBO J.* **19**:542-549.
 165. Sharma, V., A. Arockiasamy, D. R. Ronning, C. G. Savva, A. Holzenburg, M. Braunstein, W. R. Jacobs, Jr., and J. C. Sacchettini. 2003. Crystal structure of *Mycobacterium tuberculosis* SecA, a preprotein translocating ATPase. *Proc.Natl.Acad.Sci.U.S.A* **100**:2243-2248.
 166. Shimizu, H., K. Nishiyama, and H. Tokuda. 1997. Expression of *gpsA* encoding biosynthetic sn-glycerol 3-phosphate dehydrogenase suppresses both the LB- phenotype of a *secB* null mutant and the cold-sensitive phenotype of a *secG* null mutant. *Mol.Microbiol.* **26**:1013-1021.
 167. Shimoike, T., Y. Akiyama, T. Baba, T. Taura, and K. Ito. 1992. SecY variants that interfere with *Escherichia coli* protein export in the presence of normal SecY. *Mol.Microbiol.* **6**:1205-1210.
 168. Shimokawa, N., H. Mori, and K. Ito. 2003. Importance of transmembrane segments in *Escherichia coli* SecY. *Mol.Genet.Genomics.* **269**:180-187.
 169. Snyder, S., V. Ramamurthy, and D. Oliver. 1997. Identification of a region of interaction between *Escherichia coli* SecA and SecY proteins. *J.Biol.Chem.* **272**:11302-11306.
 170. Spelbrink, R. E., A. Kolkman, M. Slijper, J. A. Killian, and B. de Kruijff. 2005. Detection and identification of stable oligomeric protein complexes in *Escherichia coli* inner membranes: a proteomics approach. *J.Biol.Chem.* **280**:28742-28748.

171. **Stenberg, F., P. Chovanec, S. L. Maslen, C. V. Robinson, L. L. Ilag, G. von Heijne, and D. O. Daley.** 2005. Protein complexes of the *Escherichia coli* cell envelope. *J.Biol.Chem.* **280**:34409-34419.
172. **Story, R. M., H. Li, and J. N. Abelson.** 2001. Crystal structure of a DEAD box protein from the hyperthermophile *Methanococcus jannaschii*. *Proc.Natl.Acad.Sci.U.S.A* **98**:1465-1470.
173. **Suzuki, H., K. Nishiyama, and H. Tokuda.** 1998. Coupled structure changes of SecA and SecE revealed by the synthetic lethality of the secAcsR11 and ΔsecE::kan double mutant. *Mol.Microbiol.* **29**:331-341.
174. **Swaving, J., K. H. van Wely, and A. J. M. Driessen.** 1999. Preprotein translocation by a hybrid translocase composed of *Escherichia coli* and *Bacillus subtilis* subunits. *J.Bacteriol.* **181**:7021-7027.
175. **Szyrach, G., M. Ott, N. Bonnefoy, W. Neupert, and J. M. Herrmann.** 2003. Ribosome binding to the Oxa1 complex facilitates co-translational protein insertion in mitochondria. *EMBO J.* **22**:6448-6457.
176. **Takamatsu, D., B. A. Bensing, and P. M. Sullam.** 2004. Four proteins encoded in the gspB-secY2A2 operon of *Streptococcus gordonii* mediate the intracellular glycosylation of the platelet-binding protein GspB. *J.Bacteriol.* **186**:7100-7111.
177. **Takamatsu, D., B. A. Bensing, and P. M. Sullam.** 2004. Genes in the accessory sec locus of *Streptococcus gordonii* have three functionally distinct effects on the expression of the platelet-binding protein GspB. *Mol.Microbiol.* **52**:189-203.
178. **Tam, P. C., A. P. Maillard, K. K. Chan, and F. Duong.** 2005. Investigating the SecY plug movement at the SecYEG translocation channel. *EMBO J.* **24**:3380-3388.
179. **Tani, K., H. Tokuda, and S. Mizushima.** 1990. Translocation of ProOmpA possessing an intramolecular disulfide bridge into membrane vesicles of *Escherichia coli*. Effect of membrane energization. *J.Biol.Chem.* **265**:17341-17347.
180. **Taura, T., T. Baba, Y. Akiyama, and K. Ito.** 1993. Determinants of the quantity of the stable SecY complex in the *Escherichia coli* cell. *J.Bacteriol.* **175**:7771-7775.
181. **Taura, T., T. Yoshihisa, and K. Ito.** 1997. Protein translocation functions of *Escherichia coli* SecY: *in vitro* characterization of cold-sensitive SecY mutants. *Biochimie* **79**:517-521.
182. **Tian, H., D. Boyd, and J. Beckwith.** 2000. A mutant hunt for defects in membrane protein assembly yields mutations affecting the bacterial signal recognition particle and Sec machinery. *Proc.Natl.Acad.Sci.U.S.A.* **97**:4730-4735.
183. **Tian, P. and I. Andricioaei.** 2006. Size, motion and function of the SecY translocon revealed by molecular dynamics simulations with virtual probes. *Biophys.J.* **90**:2718-2730.
184. **Traxler, B. and C. Murphy.** 1996. Insertion of the polytopic membrane protein MalF is dependent on the bacterial secretion machinery. *J.Biol.Chem.* **271**:12394-12400.
185. **Tziatzios, C., D. Schubert, M. Lotz, D. Gundogan, H. Betz, H. Schagger, W. Haase, F. Duong, and I. Collinson.** 2004. The bacterial protein-translocation complex: SecYEG dimers associate with one or two SecA molecules. *J.Mol.Biol.* **340**:513-524.
186. **Ulbrandt, N. D., E. London, and D. B. Oliver.** 1992. Deep penetration of a portion of *Escherichia coli* SecA protein into model membranes is promoted by anionic phospholipids and by partial unfolding. *J.Biol.Chem.* **267**:15184-15192.
187. **Ulbrandt, N. D., J. A. Newitt, and H. D. Bernstein.** 1997. The *E. coli* signal recognition particle is required for the insertion of a subset of inner membrane proteins. *Cell* **88**:187-196.

188. Urbanus, M. L., P. A. Scotti, L. Froderberg, A. Saaf, J. W. de Gier, J. Brunner, J. C. Samuelson, R. E. Dalbey, B. Oudega, and J. Luirink. 2001. Sec-dependent membrane protein insertion: sequential interaction of nascent FtsQ with SecY and YidC. *EMBO Rep.* **2**:524-529.
189. Van den Berg, B., W. M. Jr. Clemons, I. Collinson, Y. Modis, E. Hartmann, S. C. Harrison, and T. A. Rapoport. 2004. X-ray structure of a protein-conducting channel. *Nature* **427**:36-44.
190. Van der Does, C., J. de Keyzer, M. van der Laan, and A. J. M. Driessen. 2003. Reconstitution of purified bacterial preprotein translocase in liposomes. *Methods Enzymol.* **372**:86-98.
191. Van der Does, C., T. den Blaauwen, J. G. de Wit, E. H. Manting, N. A. Groot, P. Fekkes, and A. J. M. Driessen. 1996. SecA is an intrinsic subunit of the *Escherichia coli* preprotein translocase and exposes its carboxyl terminus to the periplasm. *Mol. Microbiol.* **22**:619-629.
192. Van der Does, C., E. H. Manting, A. Kaufmann, M. Lutz, and A. J. M. Driessen. 1998. Interaction between SecA and SecYEG in micellar solution and formation of the membrane-inserted state. *Biochemistry* **37**:201-210.
193. Van der Sluis, E. O., N. Nouwen, and A. J. M. Driessen. 2002. SecY-SecY and SecY-SecG contacts revealed by site-specific crosslinking. *FEBS Lett.* **527**:159.
194. Van der Sluis, E. O. and A. J. M. Driessen. 2006. Stepwise evolution of the Sec machinery in Proteobacteria. *Trends Microbiol.* **14**:105-108.
195. Van der Sluis, E. O., E. van der Vries, G. Berrelkamp, N. Nouwen, and A. J. M. Driessen. 2006. Topologically fixed SecG is fully functional. *J. Bacteriol.* **188**:1188-1190.
196. Van der Wolk, J. P., J. G. de Wit, and A. J. M. Driessen. 1997. The catalytic cycle of the *Escherichia coli* SecA ATPase comprises two distinct preprotein translocation events. *EMBO J.* **16**:7297-7304.
197. Van der Wolk, J. P., P. Fekkes, A. Boorsma, J. L. Huie, T. J. Silhavy, and A. J. M. Driessen. 1998. PrlA4 prevents the rejection of signal sequence defective preproteins by stabilizing the SecA-SecY interaction during the initiation of translocation. *EMBO J.* **17**:3631-3639.
198. Van der Wolk, J. P., M. Klose, E. Breukink, R. A. Demel, B. de Kruijff, R. Freudl, and A. J. M. Driessen. 1993. Characterization of a *Bacillus subtilis* SecA mutant protein deficient in translocation ATPase and release from the membrane. *Mol. Microbiol.* **8**:31-42.
199. Van der Laan, M., N. Nouwen, and A. J. M. Driessen. 2004. SecYEG proteoliposomes catalyze the $\Delta\psi$ -dependent membrane insertion of FtsQ. *J. Biol. Chem.* **279**:1659-1664.
200. Van Voorst, F., C. van der Does, J. Brunner, A. J. M. Driessen, and B. de Kruijff. 1998. Translocase-bound SecA is largely shielded from the phospholipid acyl chains. *Biochemistry* **37**:12261-12268.
201. Van Wely, K. H. M., J. Swaving, and A. J. M. Driessen. 1998. Translocation of the precursor of alpha-amylase into *Bacillus subtilis* membrane vesicles. *Eur. J. Biochem.* **255**:690-697.
202. Veenendaal, A. K. J., C. van der Does, and A. J. M. Driessen. 2004. The protein-conducting channel SecYEG. *Biochim. Biophys. Acta* **1694**:81-95.
203. Veenendaal, A. K. J., C. van der Does, and A. J. M. Driessen. 2001. Mapping the sites of interaction between SecY and SecE by cysteine scanning mutagenesis. *J. Biol. Chem.* **276**:32559-32566.
204. Veenendaal, A. K. J., C. van der Does, and A. J. M. Driessen. 2002. The core of the bacterial translocase harbors a tilted transmembrane segment 3 of SecE. *J. Biol. Chem.* **277**:36640-36645.

-
205. **Velankar, S. S., P. Soutanas, M. S. Dillingham, H. S. Subramanya, and D. B. Wigley.** 1999. Crystal structures of complexes of PcrA DNA helicase with a DNA substrate indicate an inchworm mechanism. *Cell*. **97**:75-84.
206. **Vrontou, E. and A. Economou.** 2004. Structure and function of SecA, the preprotein translocase nanomotor. *Biochim.Biophys.Acta*. **1694**:67-80.
207. **Vrontou, E., S. Karamanou, C. Baud, G. Sianidis, and A. Economou.** 2004. Global co-ordination of protein translocation by the SecA IRA1 switch. *J.Biol.Chem.* **279**:22490-22497
208. **Wang, L., A. Miller, S. L. Rusch, and D. A. Kendall.** 2004. Demonstration of a specific *Escherichia coli* SecY-signal peptide interaction. *Biochemistry*. **43**:13185-13192.
209. **Weiss, J. B., P. H. Ray, and P. J. Bassford, Jr.** 1988. Purified SecB protein of *Escherichia coli* retards folding and promotes membrane translocation of the maltose-binding protein *in vitro*. *Proc.Natl.Acad.Sci.U.S.A.* **85**:8978-8982.
210. **Wild, K., M. Halic, I. Sinning, and R. Beckmann.** 2004. SRP meets the ribosome. *Nat.Struct.Mol.Biol.* **11**:1049-1053.
211. **Woodbury, R. L., S. J. Hardy, and L. L. Randall.** 2002. Complex behavior in solution of homodimeric SecA. *Protein Sci.* **11**:875-882.
212. **Woolf, P. J. and J. J. Linderman.** 2003. Self organization of membrane proteins via dimerization. *Biophys.Chem.* **104**:217-227.
213. **Xu, Z., J. D. Knafels, and K. Yoshino.** 2000. Crystal structure of the bacterial protein export chaperone secB. *Nat.Struct.Biol.* **7**:1172-1177.
214. **Yahr, T. L. and W. T. Wickner .** 2000. Evaluating the oligomeric state of SecYEG in preprotein translocase. *EMBO J.* **19**:4393-4401.
215. **Zhou, J. and Z. Xu.** 2003. Structural determinants of SecB recognition by SecA in bacterial protein translocation. *Nat.Struct.Biol.* **10**:942-947.
216. **Zhou, J. and Z. Xu.** 2005. The structural view of bacterial translocation-specific chaperone SecB: implications for function. *Mol.Microbiol.* **58**:349-357.
217. **Zito, C. R. and D. Oliver.** 2003. Two-stage binding of SecA to the bacterial translocon regulates ribosome-translocon interaction. *J.Biol.Chem.* **278**:40640-40646.
